



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			



Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

Entrez PubMed

☐ 1: Arch Biochem Biophys 1986 Jan;244(1):161-8[Related Articles, Links](#)

Role of plasma and serum proteases in the degradation of elastin.

Romero N, Tinker D, Hyde D, Rucker RB.

PubMed Services

Accelerated proteolysis of tropoelastin and elastin occurs in the major arteries of chicks fed copper-deficient diets. Signs of elastin degradation are not obvious in normal arteries of copper-supplemented chicks. It is proposed that the sources of proteases that effect elastin degradation are from plasma and serum. Both calcium-dependent proteases and kallikrein were effective in degrading tropoelastin and partially crosslinked insoluble elastin into peptides similar to those detected in aortic extracts from copper-deficient chicks. As dietary copper deficiency progresses it is also possible to detect elastin peptides in plasma.

Related Resources

PMID: 3511842 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

VMD 6.0005 10 JUL 17